INSIDE alcohol and judgment

parental influence on use of alcohol

proven skills to prevent underage drinking

how alcohol affects the teenage brain

Underage

Think underage drinking doesn't affect a teen's brain? Think again.

T'S A FACT. New research shows alcohol affects a teenager's developing brain differently than an adult's. Memory, learning and impulse control can be impaired seriously. The risk of addiction goes up dramatically. So talk to your kids about the dangers of alcohol and set clear rules about no alcohol use:

Know where your kids are and what they're doing.

Visit the new website for parents with information to help prevent underage drinking





C. L. "BUTCH" OTTER

We need your help! You have more power over the choices your children make than you may realize. Studies show parental disapproval is the number one reason kids choose not to

The disturbing fact is that underage drinking is a serious problem in Idaho. Heavy binge drinking begins as early as the sixth grade, and new scientific evidence proves underage drinking can cause permanent damage to a teen's rapidly developing brain. As a result, the need for parents to stay involved in their children's lives has never been greater.

The most effective parenting techniques are among the most simple, including:

- Setting clear rules about no underage drinking
- Knowing where your children are and whom they are with
- Knowing your children's friends
- Ensuring your children's environments are alcohol-free
- Having daily, positive communication and interaction with your children

The State of Idaho is urging parents to join us in fighting this severe threat to our children by educating themselves about the dangers of underage drinking and the proven skills to prevent it. Visit http://www.odp.idaho.gov today for more information. Our children deserve to have the brightest possible future we can give them, so start talking before they start drinking.

We wish you and your family all the best.

Sincerely,

Governor of Idaho

STATE CAPITOL • BOISE, IDAHO 83720 • (208) 334-2100

Teen brain development and



Most parents talk to their kids about drinking two years too late. Age 8 is not too early.

The brain's hippocampus (responsible for learning and memory) can be 10% smaller in underage drinkers.

Alcohol affects a teen brain differently from an adult brain.


It can actually cause serious damage to the still-developing adolescent brain (10-21 years).

EW RESEARCH SHOWS that alcohol affects a developing teen brain differently from an adult brain. "The brain goes through dynamic changes during adolescence, and alcohol can seriously damage longand short-term growth processes" (American Medical Association, 2003).

Alcohol use may impair memory, learning, decision-making and impulse control; and it greatly increases the risk of addiction. In addition, alcohol use can cause young people to develop social problems, have poor judgment, get into trouble, do poorly in school and experience failure in achieving lifelong goals.

To compound this problem, research shows that parents generally underestimate the extent of teen drinking and its negative consequences. Thirty-one percent of youths who said they had been drunk in the past year were said by their parents to be non-drinkers. Others may view underage drinking as inevitable, but it isn't. To be alcohol-free, a teen needs the active involvement and help of a parent. All children need help from their parents to guide them through their teenage years. Staying bonded to parents is critical to the process of teenagers emerging healthy, safe and alcohol-free.

Increased risk of





HE BRAIN REWARDS positive actions with feelings of pleasure so we want to repeat them. We remember pleasure from dopamine, a "feel-good" brain chemical, or neurotransmitter, which connects the pleasure to the thing we enjoyed. Alcohol affects our brain's pleasure-reward system by pretending to be a neurotransmitter. It tricks the

If your family has a history of alcoholism. your children need to know that they are at a greater risk for problem drinking.

brain into generating pleasure-reward feelings from a harmful chemical instead of a real experience.


If a teen continues drinking, the brain changes and adapts to the presence of alcohol and soon the teen needs more and more alcohol to create the same amount of pleasure.

If a teen still continues drinking, he or she will begin to feel a craving for it and feel uncomfortable – sometimes even extreme discomfort – without it. He or she becomes addicted. Getting their next drink becomes more important than family, grades or even sports.

Because the teen brain produces an abundance of dopamine, it can go rapidly from liking, to wanting, to needing alcohol, programming it for alcoholism. Alcohol can also damage the brain's ability to sense pleasure from normal, healthy things and experiences - leaving a young person feeling "flat" about things he/ she previously enjoyed.

A person who starts drinking at the legal age of 21 has only a 7 percent chance of becoming addicted.

Children who begin drinking at age 13 have a 45 percent chance of becoming alcohol dependent.



Parental

STARTTALKING

BEFORE THEY

START DRINKING

Sharing values and family history regarding alcohol will create an environment of trust and understanding.

If parents drink, they should:

Be clear that they do not want their children to drink alcohol until they are 21 years old and then only in moderation.

Tell their children that some people should not drink alcoholic beverages at all. These include:

- Children and adolescents.
- Individuals of any age who have a problem restricting their drinking to moderate levels.
- Women who are pregnant.
- People who plan to drive or take part in activities that require attention or skill
- People using prescription and over-the-counter medications.

If parents don't drink, they should:

- Explain their reasons for not drinking, whether they are religious, health-related or due to family history.
- Encourage children to talk about concerns and questions about drinking.
- Be clear that they do not want their children to drink.
- Explain that when the children are 21, if they should decide to drink, they should do so in moderation.

influence on alcohol use

DUCATION ALONE is not enough to deter teens from drinking as they enter difficult social transitions to adulthood, because there are many pressures and opportunities to drink. Parents who are actively involved can have a powerful influence on their child's decision to remain alcohol-free.

Teen perceptions of parental disapproval are great deterrents: What parents may not realize is that children say parental disapproval of underage drinking is the key reason they have chosen not to drink.

Research indicates that children are less likely to drink when their parents are involved in their lives and when they and their parents report feeling close to each other. Family conflict and lack of bonding are associated with increased risk of drinking. Mixed messages, and unclear rules and expectations also leave children more vulnerable to underage drinking.

Binge drinking now begins as early as elementary school, and parents are often unaware of their child's use of alcohol.

In fact, a national survey *found 31 percent of kids* who said they had been drunk in the past year had parents who believed their children to be non-drinkers.

How parents can help their children to remain

ALCOHOL-FREE



Learn and explain the risks of underage drinking. Emphasize that drinking alcohol is not a "rite of passage" but a dangerous drug for a teen brain.



Talk early and often. In Idaho, surveys indicate some youths binge drink in the sixth grade, and a few may start even earlier.



Set clear rules. Set clear rules about no alcohol use. Be specific: "Absolutely no underage drinking in our family."



Know your children's friends. Get to know your children's friends and their parents. Help them choose friends who support your family rules.



Monitor children's activities. Always know where your children are, whom they are with and what they are doing. For example: "If alcohol is at a party, call me and I'll come and get you.'



Make alcohol unavailable. Ensure that alcohol is not available to your child at home or from others when your child is away.



Be involved. Develop close bonding experiences and have daily positive interactions with your



Stay in contact. Studies show children

are more likely to drink between the hours of 3:00-6:00 PM, when unsupervised by parents. Give your kids a call.

UNDERAGE DRINKING

Underage

Some Idaho kids even begin drinking in the sixth grade. Talk to your children about not drinking early. Begin talking before age 8.

About the brain:

The **brain** is the major organ of the central nervous system and the control center for all the body's voluntary and involuntary activities. It is responsible for everything we think, feel, see, say and do. The **brainstem** controls vital body functions, such

as breathing and digestion. The **cerebellum's** main

functions are the maintenance of posture and the coordination of body movements. The

cerebrum, which consists of the right and left cerebral hemispheres, is the site of most conscious

and intelligent activities.

The brain's hippocampus (responsible for learning and memory) can be 10% smaller in

underage drinkers.

Cerebral cortex

brain research

and the effects of alcohol

the brain is highly plastic and shaped by experience.... Alcohol appears to interfere with the changes in circuitry that occur during learning." Dr. Aaron White, Duke University

SAGITTAL SECTION

"Teens, through their choices and actions, have the power to direct the development of their own brains.'

"... it has become clear that, during adolescence."

Dr. Jay Giedd, Chief of Brain Imaging, NATIONAL INSTITUTES OF HEALTH

"The use of alcohol, by itself or with other drugs, can retard the normal growth and development of young people."

> AMERICAN ACADEMY OF PEDIATRICS, "ALCOHOL: YOUR CHILD AND DRUGS"

"In sometimes subtle and sometimes dramatic ways, underage alcohol use can sidetrack the trajectory of a child's life - or end it."

Dendrites

Soma

termina

Sends an electrical

impulse down the

axon to the axon

THE U.S. SURGEON GENERALS"S CALL TO ACTION, 2007

LCOHOL AFFECTS a teen brain differently than it affects a mature adult brain. The brain goes through rapid development and "wiring" changes during the ages of 12 through 21. Teen alcohol use can damage this brain wiring, which is essential to become a mature. thoughtful, responsible adult.

THE BRAIN

Alcohol can damage two kev brain areas:

"The **prefrontal** area (responsible for thinking, planning, good judgment, decision-making and impulse control) undergoes the most change during adolescence. Researchers found that adolescent drinking could cause severe changes in this area ... which plays an important role in forming adult personality and behavior. Damage from alcohol at this time can be long-term and irreversible."

"The **hippocampus** (involved in learning and memory) suffers the worst alcohol-related brain damage in teens. Those who had been drinking more and for longer had significantly smaller hippocampi (10 percent). In addition, short-term or moderate drinking impairs learning and memory far more in youths than adults. Frequent drinkers may never be able to catch up in adulthood, since alcohol inhibits systems crucial for storing new information."

American Medical Association Fact Sheet, 2003

Occipital lobe Frontal lobe Cerebellum Olfactory bulb Pituitary gland *Hippocampus* **PREFRONTAL** AREA

Hypothalamu

Pineal gland

Alcohol acts like a computer virus in the brain by:


__Slowing or shutting down brain activity.

Deleting or distorting neural messages.

Damaging neuro-connections.

Hindering formation of mature brain wiring.

- Rewiring a brain's pleasure-reward system, which can lead to alcohol dependence/addiction.



Brain plasticity and grow new dendrites as a result of new learning and experiences.

Neural plasticity refers to the ability of circuitry in the brain to physically change

During peaks of plasticity, the adolescent brain is "wiring" two • important brain areas: the **prefrontal cortex** (responsible for • planning, decision-making, good judgment and impulse control) and the • **hippocampus** (involved in memory and learning). During this period of development, the brain must make the key neural connections to wire itself to become a responsible, thoughtful adult. Alcohol, which acts as a depressant, slows down brain activity and hinders development.

How neurons communicate

- Axon terminals Axon terminals give Receive a message from Axon (trunk) the body and send it to Dendrite receptors pick up the

> Our brain is more complex than the world's most powerful computer. It is responsible for everything we think, see, hear, feel, smell or do. It even creates and directs all of our emotions. The brain is divided into different areas that direct different parts of our body. Like a complex computer, all the different parts of the brain can work at the same time - like parallel processing.

message and forward it to the soma

It does this through more than 100 billion brain cells called neurons. A neuron looks sort of like a tree, with "branches" called dendrites, a "trunk" called an axon, and "roots" called axon terminals. The tip of each "root" contains tiny sacks of powerful chemicals called neurotransmitters. At the top of the "trunk" is a tiny electrical

The brain neurons communicate by sending electrical and chemical messages from the "roots" of one neuron to the "branches" of another. If a thought or action is repeated often, the "roots" of one neuron send more chemical, and the receiving neuron makes more "branches" to receive it. The neural connection is strengthened until it begins to look like a bushy tree instead of a spindly tree. It becomes a dominant neural pathway

40 percent of our neurons are "wired" at birth. They perform automatic functions such as breathing, heart and lung functions, digestion, etc. The other 60 percent are waiting to be stimulated by our learning and experiences to make connections or "wiring." When we learn new things or do new activities, new "NEURAL CONNECTIONS" are made in our brain. This is referred to as "wiring" our brain. It is like loading new software into a computer so it can do more things. The more neural connections we make, the smarter and more capable we become.

Alcohol and judgment

The teenage brain is still developing. Alcohol can impair the parts of the brain that control the following:

This includes the abi<mark>l</mark>ity to talk, drive and process

Motor coordination. Impulse control. Drinking lowers Memory. Impaired inhibitions and increases the chances that a person will do something that they will regret when they are sober.

recollection and even blackouts can occur when too much alcohol has

Judgment and decisionmaking capacity. Drinking may lead young people to engage in risky behaviors that can result in illness, injury and

Risks associated

with underage

drinking

Alcohol use among children is strongly correlated with violence, poor academic performance and other harmful behaviors. Education alone will not keep your kids from using alcohol because there are constant pressures and opportunities to drink.

ALCOHOL POISONING

Most kids have not vet developed the "cut-off" switch that makes them go to sleep or pass out from too much drinking. They can consume dangerous amounts of alcohol before they realize it's too late. This can result in alcohol poisoning. which can cause difficulty breathing, unconsciousness and death. Binge drinking can and does kill killing as many young people as all other drugs combined. If a young person ever passes out from drinking, 911 should be called for immediate medical

attention.

Further, the areas of the brain that encourage impulsivity and risk-taking develop early in a teen, while the areas that improve self-control don't develop until the very late teens or early twenties. You must stay actively involved as a parent in bonding with your teen, setting boundaries and monitoring to help your teen remain alcohol-free.

Violence

Children who start drinking before age 15 are 12 times more likely to be injured while under the influence of alcohol and 10 times more likely to be in a fight after drinking, compared with those who wait to drink until they are 21.

School

Teens who use alcohol have higher rates of academic problems and poor performance than non-drinkers. Among eighth-graders, higher truancy rates are associated with greater rates of alcohol use.

Promiscuity

Alcohol use by teens is a strong predictor of unprotected sexual activity and unwanted sexual advances.

Traffic deaths

Auto crashes are the number one killer of teens and more than one third of teen traffic deaths are alcohol-related. In the past year 28 Idaho children and young adults died in crashes involving drinking drivers. Thirty per cent of Idaho high school students reported riding with a drinking driver in the past month.

Illicit drug use

More than 67 percent of young people who start drinking before the age of 15 will try an illicit drug. Children who drink are 7.5 times more likely to use any illicit drug, more than 22 times more likely to use marijuana and 50 times more likely to use cocaine than children who never drink.



Stay in contact

Be aware that studies show kids are more at risk for alcohol between the hours of 3 p.m. and 6 p.m., while many parents are still at work.

Find ways to check on your kids when you are not around, either with phone calls, text messaging or through a neighbor dropping by.



Help your child to choose friends wisely

Get to know your child's friends and their parents. Discuss your no-alcohol rule with the parents and enlist their support. Ensure that teens and their friends have planned activities with appropriate adult supervision.

> lf your child's friends drink encourage them to stop.

Friends and peers

The single most predictive risk for underage drinking is if your child's peers drink. Encourage your kids to choose friends who support your family values and no-alcohol rules.

Students with high-refusal-assertiveness skills are less likely to drink underage. Decide good ways to say "no" and practice them often in role-play situations. Some ideas are:

- "No thanks. Drinking is not my thing."
- "No thanks. I need all the brain cells I've got."
- "No thanks. I've only got one brain. Why would I want to
- "No thanks. Drinking before your brain is developed can dumb yourself down."

Possessing, purchasing or drinking alcohol before age 21 is illegal.

The law

It is illegal for anyone to furnish or supply alcohol to a minor — punishable by up to a \$1,000 fine and 12 months in jail.



Peers

Children often think that other people their age are drinking regularly, but most are not.

8 UNDERAGE DRINKING

Encouraging news:

Central to this media campaign to eliminate underage drinking is the encouraging news that studies show parent disapproval is the **No. 1** reason children choose not to drink alcohol. Parents empowered can trump peer pressure.

The most effective parenting techniques are among the most simple, including:

- Set clear rules about no underage drinking
- Know where your children are and whom they are with
- Know your children's friends
- Ensure your children's social environments are alcohol-free
- Have daily, positive communication and interaction with your children
- **E**at dinner together

Research shows
teens who regularly
eat as a family
(5-7 times per week)
are 33 percent less
likely to use alcohol.



Most parents don't realize they are the **No. 1** influence in their children's lives.

When parents were asked to draw the greatest influences on their teens, they placed themselves last.



When teens were asked to draw the major influences in their lives, they placed their parents first, before peers and activities.



children make than you may realize. **Become a parent advocate.**

We all can do many things to support our children to make good choices about alcohol. In fact, every day citizens in communities across Idaho are working together to reduce the availability of and access to alcohol by youth, to reduce the opportunities for youth to drink, and to provide an environment where the demand for alcohol is lessened. Some of the issues that community groups are working on include:

You have more power over the choices your

- Alcohol at underage parties: When adults allow minors to drink, or provide alcohol to minors, they are committing a misdemeanor. In addition, teenagers interpret it as being given blanket permission to drink even when adults aren't around. Adults should not allow underage drinking in their presence, in their homes, or on their property under any circumstances.
- Alcohol advertising: American are subjected to \$4 billion in alcohol marketing each year in the form of advertising and product placements on youth-oriented television and radio shows, internet sites and billboards.
- **Alcohol and Minors:** Do not purchase *or* provide alcohol for people under 21.
- Alcoholic energy drinks: A new wave of "energy" drinks have hit the market that both contain alcohol and stimulants (e.g. caffeine) and are marketed directly at teenagers. It can be very difficult to distinguish the beverages containing alcohol and the percentage of alcohol in the drinks can be as high as 11%.
- **Checking of IDs:** Every time alcohol is purchased, the sales clerk should check the ID of the purchaser.
- **Theft of alcohol:** The theft of alcohol by young people is a serious problem and "beer runs" are a common occurrence. Sometimes the easiest place to get alcohol is right next to the milk in the refrigerator, so parents need to ensure that any alcohol in the home is inaccessible.

Idaho Laws and Penalties

Penalties of providing alcohol to underage youth:

A person who sells or furnishes alcohol to an underage youth will face fines between \$500 and \$1,000 plus up to one year in jail on the first offense, more on the second.

Penalties/Laws on underage alcohol use or possession:

- It is illegal for those under 21 to drink, buy, possess, transport or have open alcohol. Penalties for a minor in possession of alcohol: 1st offense can be a fine up to \$1,000 and up to one year driver's license suspension.
- It is illegal to have a fake ID to purchase alcohol. Penalty is a general misdemeanor: up to \$1,000 fine, 6 months in jail.
- Those who refuse alcohol tests when driving a motor vehicle and stopped by a law enforcement officer could lose their license and get points on their driving record.
- In Idaho, people under age 21 who are found with a blood alcohol level of .02 .07 face charges of Underage Driving Under the Influence with a 1st offense maximum penalty of a \$1,000 fine and a one year driver's license suspension.

Resources

If you believe your child or teen may be using drugs or alcohol and you want to find help, you can call:

- A counselor at your child's school. The child may be referred to a student assistance program with education, counseling, and support groups. Counselors may determine if there is a need for a complete assessment to determine possible treatment needs.
- 2. Your family physician or clergy.
- 3. Your county juvenile probation office may offer suggestions or classes to youth.

If treatment is needed, financial assistance may be available, depending on family income and existing insurance. Call BPA at 1-800-922-3406 to determine eligibility and for guidance in locating services.

Idaho 211 Careline

Lists low cost or free health and human services in your community.

1-800-926-5288 or 2-1-1 or www.idahocareline.org

Health and Welfare Substance Abuse Programs

http://SubstanceAbuse.idaho.gov

Children's Mental Health Councils

www.helpingldahoFamilies.com for information regarding children's behavioral, emotional, or mental health concerns.

Substance Abuse Prevention Services

www.PreventionIdaho.net includes classes for children and parenting classes.

Idaho Federation of Families

www.idahofederation.org focuses on helping families meet the needs of youth with mental, emotional, or behavioral disorders.

Treasure Valley Alcohol Drug Coalition

www.tvadc.org

For More Information

Underage Drinking Enforcement Training Center

Idaho Regional Alcohol and Drug Awareness Resource Center (RADAR)

http://hs.boisestate.edu/RADAR

Idaho College Health Coalition

www.boisestate.edu/healthservices/ICHC

Idaho State Dept. of Education

www.sde.state.id.us/Safe

Surgeon General's Call to Action to Prevent and Reduce Underage Drinking

www.surgeongeneral.gov or www.stopalcoholabuse.gov

Leadership to Keep Children Alcohol Free www.alcoholfreechildren.org

Idaho Office of Drug Policy

http://www.odp.idaho.gov

Credits

This publication was made possible through the Idaho Office of Drug Policy, the Enforcing Underage Drinking Laws Partnership, Department of Juvenile Corrections, Treasure Valley Alcohol Drug Coalition, Jacksons Food Stores, Idaho PTA, Meridian Mayor's Anti-Drug Coalition, Idaho Department of Health and Welfare, and grant #2006-AH-FX-0009 from the Office of Juvenile Justice and Delinquency Prevention.

11

